

Basic Information

Product Name	Anti-BCL2 Antibody		
Gene Name	BCL2		
Source	Rabbit		
Clonality	Polyclonal		
Isotype	IgG		
Species Reactivity	human		
Tested Application	WB, ICC/IF, FCM, ELISA		
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.		
Immunogen	E. coli-derived human Bcl-2 recombinant protein (Position: Q118-E165).		
Concentration	500 ug/ml		
Purification	Immunogen affinity purified.		
Observed MW	26 kDa		
Dilution Ratios	Western blot (WB):	1:500-2000	
	Immunocytochemistry/Immunofluorescence (ICC/IF):	1:50-400	
	Flow Cytometry (Fixed):	1:50-200	
	Enzyme linked immunosorbent assay (ELISA):	1:100-1000	

Storage

12 months from date of receipt, -20°C as supplied.


Background Information

Immunoreactive BCL2 protein in the neoplastic cells of almost all follicular lymphomas whereas no BCL2 protein was detected in follicles affected by nonneoplastic processes or in normal lymphoid tissue. Every tumor with molecular-genetic evidence of t(14;18) translocation expressed detectable levels of BCL2 protein, regardless of whether the breakpoint was located in or at a distance from the BCL2 gene. Overexpression of BCL2 blocks the apoptotic death of a pro-B-lymphocyte cell line.

Reference

Anti-BCL2 Antibody被引用在156文献中。

Selected Validation Data



Western blot analysis of BCL2 using anti-BCL2 antibody (A00040-2). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Jurkat whole cell lysates,

Lane 2: human HL-60 whole cell lysates,

Lane 3: human THP-1 whole cell lysates.

After electrophoresis, proteins were transferred to a membrane.

Then the membrane was incubated with rabbit anti-BCL2 antigen

affinity purified polyclonal antibody (A00040-2) at a dilution of


1:1000 and probed with a goat anti-rabbit IgG-HRP secondary

antibody (Catalog # BA1054). The signal is developed using ECL Plus

Western Blotting Substrate (Catalog # AR1197). A specific band was

detected for BCL2 at approximately 26 kDa. The expected band size

for BCL2 is at 26 kDa.



ICC/IF analysis of BCL2 using anti-BCL2 antibody (A00040-2) and anti-Alpha Tubulin antibody (M03989-3).

BCL2 was detected in an immunocytochemical section of U2OS cells.


The section was incubated with rabbit anti-BCL2 Antibody

(A00040-2) at a dilution of 1:100. Cy3-conjugated Anti-rabbit IgG

Secondary Antibody (red)(Catalog#BA1032) and Fluoro488-

conjugated Anti-mouse IgG Secondary Antibody (Green) (Catalog #

BA1126) were used as secondary antibody.



Flow Cytometry analysis of U2OS cells using anti-BCL2 antibody (A00040-2).

Overlay histogram showing U2OS cells stained with A00040-2 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-BCL2 Antibody (A00040-2) at 1:100 dilution for 30 min at 20°C. Fluoro488 conjugated goat anti-rabbit IgG (BA1127) was used as secondary antibody at 1:100 dilution for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG at 1:100 dilution used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.