

Basic Information

Product Name	Anti-iNOS/NOS2 Antibody	
Gene Name	NOS2	
Source	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Species Reactivity	mouse	
Tested Application	WB, FCM, ELISA	
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.	
Immunogen	E.coli-derived mouse iNOS/Nos2 recombinant protein (Position: I40-L1099).	
Concentration	500 ug/ml	
Purification	Immunogen affinity purified.	
Observed MW	130 kDa	
Dilution Ratios	Western blot (WB):	1:500-2000
	Flow Cytometry (Fixed):	1:50-200
	Enzyme linked immunosorbent assay (ELISA):	1:100-1000

Storage

12 months from date of receipt, -20°C as supplied.


Background Information

Nitric oxide synthase, inducible is an enzyme that in humans is encoded by the NOS2 gene. Nitric oxide (NO) is a messenger molecule with diverse functions throughout the body. In the brain and peripheral nervous system, NO displays many properties of a neurotransmitter; it is implicated in neurotoxicity associated with stroke and neurodegenerative diseases, neural regulation of smooth muscle, including peristalsis, and penile erection. Three different NOS isoforms have been identified which fall into two distinct types, constitutive and inducible. The inducible NOS (iNOS) isoform is expressed in a variety of cell types and tissues in response to inflammatory agents and cytokines. The human iNOS (NOS2) gene is approximately 37 kb in length and consists of 26 exons and 25 introns. NOS2-derived NO is a prerequisite for cytokine signaling and function in innate immunity.

Reference

Anti-iNOS/NOS2 Antibody被引用在9文献中。

Selected Validation Data




Western blot analysis of anti-NOS2 antibody (A00368-4). The sample well of each lane was loaded with 30ug of sample under reducing conditions.

Lane 1: mouse RAW264.7(-LPS) whole cell lysates,

Lane 2: mouse RAW264.7(+LPS) whole cell lysates.

Use rabbit anti-NOS2 1:1000, probed with a goat anti-rabbit IgG-HRP secondary antibody. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog#EK1002). A specific band was detected for NOS2 at approximately 130KD. The expected band size for NOS2 is at 130KD.



Flow Cytometry analysis of Hepa1-6 cells using anti-iNOS/NOS2 antibody (A00368-4).

Overlay histogram showing Hepa1-6 cells stained with A00368-4 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-iNOS/NOS2 Antibody (A00368-4) at 1:100 dilution for 30 min at 20°C. Fluoro488 conjugated goat anti-rabbit IgG (BA1127) was used as secondary antibody at 1:100 dilution for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG at 1:100 dilution used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.