

Basic Information

Product Name	Anti-14-3-3 Epsilon/YWHAE Antibody
Gene Name	YWHAE
Source	Rabbit
Clonality	Polyclonal
Isotype	IgG
Species Reactivity	human, mouse, rat
Tested Application	WB, IHC, ICC/IF, FCM, ELISA
Contents	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.
Immunogen	E.coli-derived human YWHAE recombinant protein (Position: M1-Q255).
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	29 kDa
Dilution Ratios	Western blot (WB): 1:500-2000 Immunohistochemistry (IHC): 1:50-400 Immunocytochemistry/Immunofluorescence (ICC/IF): 1:50-400 Flow Cytometry (Fixed): 1:50-200 Enzyme linked immunosorbent assay (ELISA): 1:100-1000 (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.

Storage


12 months from date of receipt, -20°C as supplied.

Background Information

14-3-3 protein epsilon is a protein that in humans is encoded by the YWHAE gene. This gene product belongs to the 14-3-3 family of proteins which mediate signal transduction by binding to phosphoserine-containing proteins. This highly conserved protein family is found in both plants and mammals, and this protein is 100% identical to the mouse ortholog. It interacts with CDC25 phosphatases, RAF1 and IRS1 proteins, suggesting its role in diverse biochemical activities related to signal transduction, such as cell division and regulation of insulin sensitivity. It has also been implicated in the pathogenesis of small cell lung cancer. Two transcript variants, one protein-coding and the other non-

protein-coding, have been found for this gene.


Selected Validation Data



Western blot analysis of 14-3-3 Epsilon/YWHAE using anti-14-3-3 Epsilon/YWHAE antibody (A01687-4). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.


Lane 1: human hela whole cell lysates,
 Lane 2: human Jurkat whole cell lysates,
 Lane 3: human hepg2 whole cell lysates,
 Lane 4: human SH-SY5Y whole cell lysates,
 Lane 5: human HEK293 whole cell lysates,
 Lane 6: human SW620 whole cell lysates,
 Lane 7: human A549 whole cell lysates,
 Lane 8: human Raji whole cell lysates,
 Lane 9: Rat brain tissue lysates,
 Lane 10: Mouse brain tissue lysates,
 Lane 11: Mouse NIH/3T3 whole cell lysates,
 Lane 12: Mouse RAW264.7 whole cell lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-14-3-3 Epsilon/YWHAE antigen purified polyclonal antibody (A01687-4) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for 14-3-3 Epsilon/YWHAE at approximately 29 kDa. The expected band size for 14-3-3 Epsilon/YWHAE is at 29 kDa.




IHC analysis of 14-3-3 Epsilon/YWHAE using anti-14-3-3 Epsilon/YWHAE antibody (A01687-4).

14-3-3 Epsilon/YWHAE was detected in a paraffin-embedded section of rat brain tissue. Biotinylated goat anti-rabbit IgG was used as secondary antibody. The tissue section was incubated with rabbit anti-14-3-3 Epsilon/YWHAE Antibody (A01687-4) at a dilution of 1:200 and developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB (Catalog # AR1027) as the chromogen.



ICC/IF analysis of 14-3-3 Epsilon/YWHAE using anti-14-3-3 Epsilon/YWHAE antibody (A01687-4).

14-3-3 Epsilon/YWHAE was detected in an immunocytochemical section of MCF-7 cells. The section was incubated with rabbit anti-14-3-3 Epsilon/YWHAE Antibody (A01687-4) at a dilution of 1:100. Fluoro488 Conjugated Goat Anti-Rabbit IgG (Green) (Catalog # BA1127) was used as secondary antibody. The section was counterstained with DAPI (Catalog # AR1176) (Blue).



Flow Cytometry analysis of A549 cells using anti-14-3-3 Epsilon/YWHAE antibody (A01687-4).

Overlay histogram showing A549 cells stained with A01687-4 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-14-3-3 Epsilon/YWHAE Antibody (A01687-4) at 1:100 dilution for 30 min at 20°C. Fluoro488 conjugated goat anti-rabbit IgG (BA1127) was used as secondary antibody at 1:100 dilution for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG at 1:100 dilution used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

Product datasheet

Anti-14-3-3 Epsilon/YWHAE Antibody

Catalog Number: A01687-4



antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY

Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator,
East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com Phone: 027-67845390/1/2 Email: boster@boster.com