

## Basic Information

<b>Product Name</b>	Anti-iNOS/NOS2 Antibody	
<b>Gene Name</b>	NOS2	
<b>Source</b>	Rabbit	
<b>Clonality</b>	Polyclonal	
<b>Isotype</b>	IgG	
<b>Species Reactivity</b>	human, mouse, rat	
<b>Tested Application</b>	WB, IHC	
<b>Contents</b>	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.	
<b>Immunogen</b>	A synthetic peptide corresponding to a sequence at the C-terminus of human iNOS, identical to the related rat and mouse sequences.	
<b>Concentration</b>	500 ug/ml	
<b>Purification</b>	Immunogen affinity purified.	
<b>Observed MW</b>	130 kDa	
<b>Dilution Ratios</b>	Western blot (WB): Immunohistochemistry (IHC): (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	1:500-2000 1:50-400

## Storage

12 months from date of receipt, -20°C as supplied.

## Background Information


Nitric oxide synthase, inducible is an enzyme that in humans is encoded by the NOS2 gene. Nitric oxide (NO) is a messenger molecule with diverse functions throughout the body. In the brain and peripheral nervous system, NO displays many properties of a neurotransmitter; it is implicated in neurotoxicity associated with stroke and neurodegenerative diseases, neural regulation of smooth muscle, including peristalsis, and penile erection. Three different NOS isoforms have been identified which fall into two distinct types, constitutive and inducible. The inducible NOS (iNOS) isoform is expressed in a variety of cell types and tissues in response to inflammatory agents and cytokines. The human iNOS (NOS2) gene is approximately 37 kb in length and consists of 26 exons and 25 introns. NOS2-derived

NO is a prerequisite for cytokine signaling and function in innate immunity.

## Reference

Anti-iNOS/NOS2 Antibody被引用在132文献中。

## Selected Validation Data



Western blot analysis of iNOS/NOS2 using anti-iNOS/NOS2 antibody (BA0362). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: SCG whole cell lysates,

Lane 2: COLO 320 whole cell lysates,

Lane 3: MCF-7 whole cell lysates.

After electrophoresis, proteins were transferred to a membrane.


Then the membrane was incubated with rabbit anti-iNOS/NOS2

antigen affinity purified polyclonal antibody (BA0362) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary

antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was

detected for iNOS/NOS2 at approximately 130 kDa. The expected

band size for iNOS/NOS2 is at 131 kDa.



IHC analysis of iNOS/NOS2 using anti-iNOS/NOS2 antibody (BA0362). iNOS/NOS2 was detected in a paraffin-embedded section of human intestine tissue. Biotinylated goat anti-rabbit IgG was used as secondary antibody. The tissue section was incubated with rabbit anti-iNOS/NOS2 Antibody (BA0362) at a dilution of 1:200 and developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB (Catalog # AR1027) as the chromogen.