

Basic Information

Product Name	Anti-Nestin/NES Antibody	
Gene Name	NES	
Source	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Species Reactivity	human, mouse, rat, rabbit	
Tested Application	WB, IHC	
Contents	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.	
Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human Nestin different from the related mouse sequence by four amino acids, and from the related rat sequence by five amino acids.	
Concentration	500 ug/ml	
Purification	Immunogen affinity purified.	
Observed MW	230 kDa	
Dilution Ratios	Western blot (WB): 1:500-2000 Immunohistochemistry (IHC): 1:50-400 (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	

Storage

12 months from date of receipt, -20°C as supplied.

Background Information

Nestin is a type VI intermediate filament (IF) protein that in humans is encoded by the NES gene. This gene is mapped to 1q23.1. Distribution and expression of nestin in mitotically active cells suggests it plays a role in regulation of the assembly and disassembly of intermediate filaments, which, together with other structural proteins, participate in remodeling of the cell. The role of nestin in dynamic cells, particularly structural organization of the cell, appears strictly regulated by phosphorylation, especially its integration into heterogeneous intermediate filaments together with vimentin or α -internexin. Furthermore, nestin expression has been extensively used as a marker for central nervous

system (CNS) progenitor cells in different contexts, based on observations indicating a correlation between nestin expression and this cell type in vivo.

Reference

Anti-Nestin/NES Antibody被引用在12文献中。

Selected Validation Data



IHC analysis of Nestin/NES using anti-Nestin/NES antibody (BA1289). Nestin/NES was detected in a paraffin-embedded section of human brain cancer tissue. Biotinylated goat anti-rabbit IgG was used as secondary antibody. The tissue section was incubated with rabbit anti-Nestin/NES Antibody (BA1289) at a dilution of 1:200 and developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB (Catalog # AR1027) as the chromogen.