

Basic Information

Product Name	Anti-OLIG1 Antibody
Gene Name	OLIG1
Source	Rabbit
Clonality	Polyclonal
Isotype	IgG
Species Reactivity	human
Tested Application	WB
Contents	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.
Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human Olig1, different from the related mouse and rat sequences by four amino acids.
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	28 kDa
Dilution Ratios	Western blot (WB):1:500-2000


Storage

12 months from date of receipt, -20°C as supplied.

Background Information

Oligodendrocyte transcription factor 1(OLIG1), also known as class B basic helix-loop-helix protein 6(BHLHB6) is a protein that in humans is encoded by the OLIG1 gene. This gene is mapped to 21q22.11. This gene is a member of the oligodendrocyte lineage gene family which encodes basic helix-loop-helix transcription factors. Studies in mice show that it is coexpressed with Olig2 in neural progenitors, and has a role in the development and maturation of oligodendrocytes. OLIG1 gene promotes formation and maturation of oligodendrocytes, especially within the brain. It cooperates with OLIG2 to establish the pMN domain of the embryonic neural tube.

Selected Validation Data



Western blot analysis of OLIG1 using anti-OLIG1 antibody (BA3478). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: U87 whole cell lysates,

Lane 2: HELA whole cell lysates.

After electrophoresis, proteins were transferred to a membrane.

Then the membrane was incubated with rabbit anti-OLIG1 antigen affinity purified polyclonal antibody (BA3478) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western

Blotting Substrate (Catalog # AR1197). A specific band was detected for OLIG1 at approximately 28 kDa. The expected band size for OLIG1 is at 28 kDa.