

Basic Information

Product Name	Anti-MCM5 Antibody	
Gene Name	MCM5	
Source	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Species Reactivity	human, mouse, rat	
Tested Application	WB, IHC, ICC/IF, FCM	
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.	
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human MCM5, identical to the related rat and mouse sequences.	
Concentration	500 ug/ml	
Purification	Immunogen affinity purified.	
Observed MW	95 kDa	
Dilution Ratios	Western blot (WB):	1:500-2000
	Immunohistochemistry (IHC):	1:50-400
	Immunocytochemistry/Immunofluorescence (ICC/IF):	1:50-400
	Flow Cytometry (Fixed):	1:50-200
	(Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	

Storage


12 months from date of receipt, -20°C as supplied.

Background Information

MCM5(MINICHROMOSOME MAINTENANCE, S. CEREVISIAE, HOMOLOG OF, 5),also known as CDC46, is a protein that in humans is encoded by the MCM5 gene. MCM5 is structurally very similar to the CDC46 protein from S. cerevisiae, a protein involved in the initiation of DNA replication. And MCM5 is a member of the MCM family of chromatin-binding proteins and can interact with at least two other members of this family. The MCM5 gene is mapped on 22q12.3. The MCM5 gene contains 16 exons. The encoded protein is upregulated in the transition from the G0 to G1/S phase of the

cell cycle and may actively participate in cell cycle regulation.


Selected Validation Data




Western blot analysis of MCM5 using anti-MCM5 antibody (BA3685). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: Human Jurkat whole cell lysates,
Lane 2: Human A375 whole cell lysates,
Lane 3: Human CACO-2 whole cell lysates,
Lane 4: Human HEK293 whole cell lysates,
Lane 5: Human MCF-7 whole cell lysates,
Lane 6: Rat spleen tissue lysates,
Lane 7: Mouse thymus tissue lysates.


After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-MCM5 antigen affinity purified polyclonal antibody (BA3685) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for MCM5 at approximately 95 kDa. The expected band size for MCM5 is at 82 kDa.



IHC analysis of MCM5 using anti-MCM5 antibody (BA3685). MCM5 was detected in a paraffin-embedded section of human intestinal cancer tissue. Biotinylated goat anti-rabbit IgG was used as secondary antibody. The tissue section was incubated with rabbit anti-MCM5 Antibody (BA3685) at a dilution of 1:200 and developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB (Catalog # AR1027) as the chromogen.



ICC/IF analysis of MCM5 using anti-MCM5 antibody (BA3685). MCM5 was detected in an immunocytochemical section of U2OS cells. The section was incubated with rabbit anti-MCM5 Antibody (BA3685) at a dilution of 1:100. Fluoro594-conjugated Anti-rabbit IgG Secondary Antibody (red)(Catalog#BA1142) was used as secondary antibody. The section was counterstained with DAPI (Catalog # AR1176) (Blue).



Flow Cytometry analysis of HL-60 cells using anti-MCM5 antibody (BA3685).

Overlay histogram showing HL-60 cells stained with BA3685 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-MCM5 Antibody (BA3685) at 1:100 dilution for 30 min at 20°C. Fluoro488 conjugated goat anti-rabbit IgG (BA1127) was used as secondary antibody at 1:100 dilution for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG at 1:100 dilution used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.