

## Basic Information

<b>Product Name</b>	Anti-Survivin/BIRC5 Antibody	
<b>Gene Name</b>	BIRC5	
<b>Source</b>	Rabbit	
<b>Clonality</b>	Polyclonal	
<b>Isotype</b>	IgG	
<b>Species Reactivity</b>	human	
<b>Tested Application</b>	WB, ELISA	
<b>Contents</b>	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.	
<b>Immunogen</b>	E. coli-derived human Survivin recombinant protein(Position: M1-D142).	
<b>Concentration</b>	500 ug/ml	
<b>Purification</b>	Immunogen affinity purified.	
<b>Observed MW</b>	16 kDa	
<b>Dilution Ratios</b>	Western blot (WB):	1:500-2000
	Immunohistochemistry (IHC):	1:50-400
	Enzyme linked immunosorbent assay (ELISA):	1:100-1000
	(Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	

## Storage

12 months from date of receipt, -20°C as supplied.


## Background Information

Survivin, also called baculoviral inhibitor of apoptosis repeat-containing 5 or BIRC5, is a protein that, in humans, is encoded by the BIRC5 gene. Survivin is a member of the inhibitor of apoptosis(IAP) family. The survivin gene contains 4 exons. The survivin gene is mapped to chromosome 17q25 by pulsed field gel electrophoresis and single- and 2-color FISH. The survivin protein functions to inhibit caspase activation, thereby leading to negative regulation of apoptosis or programmed cell death. The survivin protein is expressed highly in most human tumours and fetal tissue, but is completely absent in terminally differentiated cells.

## Reference

Anti-Survivin/BIRC5 Antibody被引用在1文献中。

## Selected Validation Data



Western blot analysis of Survivin/BIRC5 using anti-Survivin/BIRC5 antibody (BA4055-2). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human PC-3 whole cell lysates,

Lane 2: human U2OS whole cell lysates,

Lane 3: human 293T whole cell lysates,

Lane 4: human A549 whole cell lysates.

After electrophoresis, proteins were transferred to a membrane.

Then the membrane was incubated with rabbit anti-Survivin/BIRC5 antigen affinity purified polyclonal antibody (BA4055-2) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for Survivin/BIRC5 at approximately 16 kDa. The expected band size for Survivin/BIRC5 is at 16 kDa.