

Basic Information

Product Name	Anti-active+pro Caspase 3/CASP3(p17) Antibody(Clone#DGI-3)
Gene Name	CASP3
Source	Rabbit
Clonality	Monoclonal
Isotype	IgG
Species Reactivity	human, mouse, rat
Tested Application	WB, IHC, IP
Contents	500 ug/ml; Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide, 0.4-0.5 mg/ml BSA and 50% glycerol.
Immunogen	A synthesized peptide derived from human active + pro Caspase 3.
Concentration	500 ug/ml
Purification	Affinity-chromatography
Observed MW	35 kDa,(cleaved)20/17/12 kDa
Dilution Ratios	Western blot (WB): 1:500-2000 Immunohistochemistry (IHC):1:50-200 ImmunoPrecipitation (IP): 1:20

Storage

12 months from date of receipt, -20°C as supplied.

Background Information


Caspase 3 is a caspase protein which interacts with Survivin, XIAP, CFLAR, Caspase 8, HCLS1, Deleted in Colorectal Cancer, TRAF3 and GroEL. This gene which is located on 4q35 encodes a protein that is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes that undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. It is the predominant caspase involved in the cleavage of amyloid-beta 4A precursor protein, which is associated with neuronal death in Alzheimer's disease. And the caspase-3 activation in heart failure sequentially cleaves SRF and generates a truncated SRF that appears to function as a dominant-negative transcription factor. Additionally, the caspase-3 influence on bone mineral density should be considered in any in vivo application of caspase-3 inhibitors to the treatment of human

disease. In erythroid precursors undergoing terminal differentiation, Hsp70 prevents active CASP3 from cleaving GATA1 and inducing apoptosis.

Reference

Anti-active+pro Caspase 3/CASP3(p17) Antibody(Clone#DGI-3)被引用在10文献中。

Selected Validation Data



Western blot analysis of anti-active + pro Caspase 3/CASP3 (p17) antibody (BM4340). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,
Lane 2: human 293T whole cell lysates,
Lane 3: human Jurkat whole cell lysates,
Lane 4: human Jurkat whole cell lysates,
Lane 5: rat liver tissue lysates,
Lane 6: rat spleen tissue lysates,
Lane 7: mouse liver tissue lysates,
Lane 8: mouse spleen tissue lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-active + pro Caspase 3/CASP3 (p17) antigen affinity purified monoclonal antibody (BM4340) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for active + pro Caspase 3/CASP3 (p17) at approximately 35 kDa,(cleaved)20/17/12 kDa. The expected band size for active + pro Caspase 3/CASP3 (p17) is at 32 kDa.

Product datasheet

Anti-active+pro Caspase 3/CASP3(p17)

Antibody(Clone#DGI-3)

Catalog Number: BM4340


BOSTER[®]

antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY

Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator,
East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com **Phone:** 027-67845390/1/2 **Email:** boster@boster.com



Immunohistochemical analysis of paraffin-embedded human stomach, using active + pro Caspase 3 Antibody .