

## Basic Information

Product Name	Anti-Histone H3 Antibody (Clone#HDE-8)	
Gene Name	H3C1/H3C2/H3C3/H3C4/H3C6/H3C7/H3C8/H3C10/H3C11/H3C12	
Source	Rabbit	
Clonality	Monoclonal	
Isotype	IgG	
Species Reactivity	human, mouse, rat	
Tested Application	WB, IHC, ICC/IF	
Contents	500 ug/ml; Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide, 0.4-0.5 mg/ml BSA and 50% glycerol.	
Immunogen	A synthesized peptide derived from human Histone H3	
Purification	Affinity-chromatography	
Observed MW	15-17 kDa	
Dilution Ratios	Western blot (WB): Immunohistochemistry (IHC): Immunocytochemistry/Immunofluorescence (ICC/IF):	1:500-2000 1:50-200 1:50-200

## Storage

12 months from date of receipt, -20°C as supplied.


## Background Information

Histones are the main constituents of the protein part of chromosomes of eukaryotic cells. They are rich in the amino acids arginine and lysine and have been greatly conserved during evolution. Histones pack the DNA into tight masses of chromatin. Two core histones of each class H2A, H2B, H3 and H4 assemble and are wrapped by 146 base pairs of DNA to form one octameric nucleosome. Histone tails undergo numerous post-translational modifications, which either directly or indirectly alter chromatin structure to facilitate transcriptional activation or repression or other nuclear processes. In addition to the genetic code, combinations of the different histone modifications reveal the so-called "histone code". Histone methylation and demethylation is dynamically regulated by respectively histone methyl transferases and histone demethylases.

## Reference

Anti-Histone H3 Antibody (Clone#HDE-8)被引用在6文献中。


## Selected Validation Data



Western blot analysis of anti-Histone H3 antibody (BM4715). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,  
 Lane 2: human Jurkat whole cell lysates,  
 Lane 3: human HepG2 whole cell lysates,  
 Lane 4: human 293T whole cell lysates,  
 Lane 5: rat heart tissue lysates,  
 Lane 6: rat liver tissue lysates,  
 Lane 7: mouse heart tissue lysates,  
 Lane 8: mouse liver tissue lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-Histone H3 antigen affinity purified monoclonal antibody (BM4715) at a dilution of 1:2000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for Histone H3 at approximately 15 kDa. The expected band size for Histone H3 is at 15 kDa.



Product datasheet

## Anti-Histone H3 Antibody

(Clone#HDE-8)

Catalog Number: **BM4715**


**BO**STER®

antibody and ELISA experts

**BOSTER BIOLOGICAL TECHNOLOGY**

Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator,  
East Lake High-Tech Development Zone, Wuhan.

**Web:** www.boster.com **Phone:** 027-67845390/1/2 **Email:** boster@boster.com



Immunofluorescent analysis of HeLa cells, using Histone H3 Antibody.