

Basic Information

Product Name	Anti-IBA1/AIF1 Antibody (Clone#AFGH-1)	
Gene Name	AIF1	
Source	Rabbit	
Clonality	Monoclonal	
Isotype	IgG	
Species Reactivity	human	
Tested Application	WB, IHC, ICC/IF, FCM	
Contents	500 ug/ml; Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide, 0.4-0.5 mg/ml BSA and 50% glycerol.	
Immunogen	A synthesized peptide derived from human IBA1	
Concentration	500 ug/ml	
Purification	Affinity-chromatography	
Observed MW	17 kDa	
Dilution Ratios	Western blot (WB):	1:500-2000
	Immunohistochemistry (IHC):	1:50-200
	Immunocytochemistry/Immunofluorescence (ICC/IF):	1:50-200
	Flow Cytometry (FCM):	1:100

Storage

12 months from date of receipt, -20°C as supplied.


Background Information

Allograft inflammatory factor 1 (AIF-1), also known as ionized calcium-binding adapter molecule 1 (IBA1), is a protein that in humans is encoded by the AIF1 gene. This gene encodes a protein that binds actin and calcium. And this gene is induced by cytokines and interferon and may promote macrophage activation and growth of vascular smooth muscle cells and T-lymphocytes. Polymorphisms in this gene may be associated with systemic sclerosis. Alternative splicing results in multiple transcript variants, but the full-length and coding nature of some of these variants is not certain.

Reference

Anti-IBA1/AIF1 Antibody (Clone#AFGH-1)被引用在8文献中。

Selected Validation Data



Western blot analysis of IBA1/AIF1 using anti-IBA1/AIF1 antibody (BM5765). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human THP-1 whole cell lysates,


Lane 2: human U937 whole cell lysates,

Lane 3: human HL-60 whole cell lysates,

Lane 4: human HEL whole cell lysates.

After electrophoresis, proteins were transferred to a membrane.

Then the membrane was incubated with rabbit anti-IBA1/AIF1 antigen affinity purified monoclonal antibody (BM5765) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for IBA1/AIF1 at approximately 17 kDa. The expected band size for IBA1/AIF1 is at 17 kDa.



IHC analysis of IBA1/AIF1 using anti-IBA1/AIF1 antibody (BM5765).

IBA1/AIF1 was detected in a paraffin-embedded section of human tonsil tissue. The tissue section was incubated with rabbit anti-IBA1/AIF1 Antibody (BM5765) at a dilution of 1:200 and developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB (Catalog # AR1027) as the chromogen.