


Basic Information

Product Name	Anti-Zebrafish ABCC12 Antibody	
Gene Name	abcc12	
Source	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Species Reactivity	zebrafish	
Tested Application	IF, IHC	
Contents	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.	
Concentration	500 ug/ml	
Purification	Immunogen affinity purified.	
Dilution Ratios	Immunohistochemistry (IHC): 1:50-400 Immunofluorescence (IF): 1:50-400 (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	


Storage

12 months from date of receipt, -20°C as supplied.

Selected Validation Data



IF analysis of ABCC12 using anti-ABCC12 antibody (DZ10852). ABCC12 was detected in a paraffin-embedded section of zebrafish embryo tissue. The tissue section was incubated with rabbit anti-ABCC12 Antibody (DZ10852) at a dilution of 1:100. Fluoro488-conjugated Anti-rabbit IgG Secondary Antibody (green) (Catalog # BA1127) was used as secondary antibody.



IHC analysis of ABCC12 using anti-ABCC12 antibody (DZ10852). ABCC12 was detected in a paraffin-embedded section of zebrafish colon tissue. The tissue section was incubated with rabbit anti-ABCC12 Antibody (DZ10852) at a dilution of 1:200 and developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB (Catalog # AR1027) as the chromogen.