

Basic Information

Product Name	Anti-Ki67/MKI67 Antibody (Clone#5E12)	
Gene Name	MKI67	
Source	Mouse	
Clonality	Monoclonal	
Isotype	IgG2b	
Species Reactivity	human	
Tested Application	IHC, IF, ICC/IF, FCM	
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.	
Immunogen	E. coli-derived human Ki67 recombinant protein (Position: K2860-I3256).	
Concentration	500 ug/ml	
Purification	protein G purified.	
Dilution Ratios	Immunohistochemistry (IHC):	1:50-400
	Immunocytochemistry/Immunofluorescence (ICC/IF):	1:50-400
	Immunofluorescence (IF):	1:50-400
	Flow Cytometry (Fixed):	1:50-200
	(Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.	

Storage

12 months from date of receipt, -20°C as supplied.


Background Information

Ki-67(Proliferation-related Ki-67 antigen), also known as MKI67 or KIA, is a protein that in humans is encoded by the MKI67 gene. From study of a panel of human-rodent somatic cell hybrids, it has been demonstrated that a gene involved in the expression of the MKI67 antigen is located on chromosome 10. By in situ hybridization, Fonatsch et al. (1991) regionalized the MKI67 gene to chromosome 10q25-qter. By FISH, Traut et al. (1998) mapped the mouse Mki67 gene to chromosome 7F3-F5. Antigen KI-67 is a nuclear protein that is associated with and may be necessary for cellular proliferation. Furthermore it is associated with ribosomal RNA transcription. Inactivation of antigen KI-67 leads to inhibition of ribosomal RNA synthesis.

Reference


Anti-Ki67/MKI67 Antibody (Clone#5E12)被引用在6文献中。

Selected Validation Data




IHC analysis of Ki67/MKI67 using anti-Ki67/MKI67 antibody (M00254-8).

Ki67/MKI67 was detected in a paraffin-embedded section of human tonsil tissue. Biotinylated goat anti-mouse IgG was used as secondary antibody. The tissue section was incubated with mouse anti-Ki67/MKI67 Antibody (M00254-8) at a dilution of 1:200 and developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB (Catalog # AR1027) as the chromogen.




IF analysis using anti-Ki67 antibody (M00254-8). detected in paraffin-embedded section of human tonsil cancer tissues. The tissue section were stained using the Fluoro488 conjugated Anti-mouse IgG Secondary Antibody ((green)(Catalog # BA1126) and counterstained with DAPI (blue).



ICC/IF analysis of Ki67/MKI67 using anti-Ki67/MKI67 antibody (M00254-8).

Ki67/MKI67 was detected in an immunocytochemical section of A549 cells. The section was incubated with mouse anti-Ki67/MKI67 Antibody (M00254-8) at a dilution of 1:100. Fluoro488-conjugated Anti-mouse IgG Secondary Antibody (green)(Catalog#BA1126) was used as secondary antibody. The section was counterstained with

DAPI (Catalog # AR1176) (Blue).



Flow Cytometry analysis of Jurkat cells using anti-Ki67/MKI67 antibody (M00254-8).

Overlay histogram showing Jurkat cells stained with M00254-8 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-Ki67/MKI67 Antibody (M00254-8) at 1:100 dilution for 30 min at 20°C. Fluoro488 conjugated goat anti-mouse IgG (BA1126) was used as secondary antibody at 1:100 dilution for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG at 1:100 dilution used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.