

## Basic Information

Product Name	Anti-Lamin A/C Antibody (Clone#5F3C12)
Gene Name	LMNA
Source	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Species Reactivity	human, mouse, rat
Tested Application	WB, IHC
Contents	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.
Immunogen	E.coli-derived human Lamin A/C recombinant protein (Position: Y481-Y646). Human Lamin A/C shares 90% and 92% amino acid (aa) sequence identity with mouse and rat Lamin A/C, respectively.
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	74 kDa
Dilution Ratios	Western blot (WB): 1:500-2000 Immunohistochemistry (IHC): 1:50-400 (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.

## Storage

12 months from date of receipt, -20°C as supplied.

## Background Information


Lamins are structural protein components of the nuclear lamina, a protein network underlying the inner nuclear membrane that determines nuclear shape and size. There are three types of lamins, A,B and C. The lamin A/C (LMNA) gene contains 12 exons. Alternative splicing within exon 10 gives rise to two different mRNAs that code for pre-lamin A and lamin C. Lamin A/C is mapped to 1q21.2-q21.3 and mutations in this gene cause a variety of human diseases including Emery-Dreifuss muscular dystrophy, dilated cardiomyopathy, and Hutchinson-Gilford progeria syndrome. Lamin A/C deficiency is thus associated with both defective nuclear mechanics and impaired mechanically activated

gene transcription.

## Reference

Anti-Lamin A/C Antibody (Clone#5F3C12)被引用在1文献中。

## Selected Validation Data



Western blot analysis of Lamin A/C using anti-Lamin A/C antibody (M00438-6). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human placenta tissue lysates,

Lane 2: human HeLa whole cell lysates,

Lane 3: human A431 whole cell lysates,

Lane 4: rat lung tissue lysates,

Lane 5: mouse lung tissue lysates,

Lane 6: mouse small intestine tissue lysates.

After electrophoresis, proteins were transferred to a membrane.


Then the membrane was incubated with mouse anti-Lamin A/C antigen affinity purified monoclonal antibody (M00438-6) at a dilution of 1:1000 and probed with a goat anti-mouse IgG-HRP secondary antibody (Catalog # BA1050). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for Lamin A/C at approximately 74 kDa.

The expected band size for Lamin A/C is at 74 kDa.

## Anti-Lamin A/C Antibody

(Clone#5F3C12)

Catalog Number: M00438-6



IHC analysis of Lamin A/C using anti-Lamin A/C antibody (M00438-6).

Lamin A/C was detected in a paraffin-embedded section of human laryngeal squamous cell carcinoma tissue. The tissue section was incubated with mouse anti-Lamin A/C Antibody (M00438-6) at a dilution of 1:200 and developed using HRP Conjugated mouse IgG Super Vision Assay Kit (Catalog # SV0001) with DAB (Catalog # AR1027) as the chromogen.