

Basic Information

Product Name	Anti-XAF1 Antibody	
Gene Name	XAF1	
Source	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Species Reactivity	human	
Tested Application	WB, ICC/IF, FCM	
Contents	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.	
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human XAF1.	
Concentration	500 ug/ml	
Observed MW	35 kDa	
Dilution Ratios	Western blot (WB):	1:500-2000
	Immunocytochemistry/Immunofluorescence (ICC/IF):	1:50-400
	Flow Cytometry (Fixed):	1:50-200


Storage

12 months from date of receipt, -20°C as supplied.

Background Information

XIAP associated factor-1, also known as XAF1, is a human gene. X-linked inhibitor of apoptosis(XIAP; MIM 300079) is a potent member of the IAP family. All members of this family possess baculoviral IAP(BIR) repeats, cysteine-rich domains of approximately 80 amino acids that bind and inhibit caspases. XAF1 antagonizes the anticaspase activity of XIAP and may be important in mediating apoptosis resistance in cancer cells. And alteration in XAF1 and XIAP RNA expression levels may lead to increased apoptotic resistance and proliferation due to unregulated XIAP function in cancer cells.

Selected Validation Data




Western blot analysis of XAF1 using anti-XAF1 antibody (PA1218).

The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Jurkat whole cell lysates.

After electrophoresis, proteins were transferred to a membrane.


Then the membrane was incubated with rabbit anti-XAF1 antigen affinity purified polyclonal antibody (PA1218) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for XAF1 at approximately 35 kDa. The expected band size for XAF1 is at 35 kDa.



ICC/IF analysis of XAF1 using anti-XAF1 antibody (PA1218).

XAF1 was detected in an immunocytochemical section of U2OS cells.

The section was incubated with rabbit anti-XAF1 Antibody (PA1218) at a dilution of 1:100. Fluoro488 Conjugated Goat Anti-Rabbit IgG (Green) (Catalog # BA1127) was used as secondary antibody.



Flow Cytometry analysis of SiHa cells using anti-XAF1 antibody (PA1218).

Overlay histogram showing SiHa cells stained with PA1218 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer.

The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-XAF1 Antibody (PA1218) at 1:100 dilution for 30 min at 20°C. Fluoro488 conjugated goat anti-rabbit IgG (BA1127) was used as secondary antibody at 1:100 dilution for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG at 1:100 dilution used under the same conditions. Unlabelled sample

Product datasheet

Anti-XAF1 Antibody

Catalog Number: **PA1218**

BOSTER[®]

antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY

Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator,
East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com **Phone:** 027-67845390/1/2 **Email:** boster@boster.com

without incubation with primary antibody and secondary antibody
(Red line) was used as a blank control.