

Basic Information

Product Name	Anti-CDK2 Antibody
Gene Name	CDK2
Source	Rabbit
Clonality	Polyclonal
Isotype	IgG
Species Reactivity	human, mouse, rat
Tested Application	WB
Contents	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Cdk2, identical to the related rat and mouse sequences.
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	30 kDa
Dilution Ratios	Western blot (WB):1:500-2000

Storage

12 months from date of receipt, -20°C as supplied.


Background Information

CDK2, Cyclin-Dependent Kinase2, is also known as P33. The CDK2 protein was highly homologous to p34(CDC2) kinase and more significantly homologous to Xenopus Eg1 kinase, suggesting that CDK2 is the human homolog of Eg1. The CDK2 gene is mapped to 12q13, the same region to which the CDK4 gene maps. Human cyclin A binds independently to 2 kinases, p34(cdc2) or p33. In adenovirus-transformed cells, the viral E1A oncoprotein seems to associate with p33/cyclin A but not with p34(cdc2)/cyclin A. The gene for p33 shares 65% sequence identity with p34(cdc2). P33(cdk2) plays a unique role in cell cycle regulation of vertebrate cells.

Reference

Anti-CDK2 Antibody 被引用在4文献中。

Selected Validation Data



Western blot analysis of CDK2 using anti-CDK2 antibody (PA1547).

The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Jurkat whole cell lysates,
Lane 2: human HepG2 whole cell lysates,
Lane 3: human U2OS whole cell lysates,
Lane 4: human K562 whole cell lysates,
Lane 5: human T-47D whole cell lysates,
Lane 6: human Caco-2 whole cell lysates,
Lane 7: human 293T whole cell lysates,
Lane 8: human PC-3 whole cell lysates.

After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-CDK2 antigen affinity purified polyclonal antibody (PA1547) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for CDK2 at approximately 30 kDa. The expected band size for CDK2 is at 30,34 kDa.