

Basic Information

Product Name	Anti-NMDAR2A/GRIN2A Antibody
Gene Name	GRIN2A
Source	Rabbit
Clonality	Polyclonal
Isotype	IgG
Species Reactivity	human, mouse, rat
Tested Application	WB
Contents	500 ug/ml antibody with PBS, 0.02% NaN ₃ , 1 mg/ml BSA and 50% glycerol.
Immunogen	E.coli-derived human NMDAR2A recombinant protein (Position: D958-R1300). Human NMDAR2A shares 89% and 90% amino acid (aa) sequence identity with mouse and rat NMDAR2A, respectively.
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	165 kDa
Dilution Ratios	Western blot (WB):1:500-2000

Storage

12 months from date of receipt, -20°C as supplied.


Background Information

GRIN2A is also known as N-methyl-D-aspartate receptor channel, subunit epsilon-1(NMDAR2A). This gene encodes a member of the glutamate-gated ion channel protein family. The encoded protein is an N-methyl-D-aspartate (NMDA) receptor subunit. NMDA receptors are both ligand-gated and voltage-dependent, and are involved in long-term potentiation, an activity-dependent increase in the efficiency of synaptic transmission thought to underlie certain kinds of memory and learning. These receptors are permeable to calcium ions, and activation results in a calcium influx into post-synaptic cells, which results in the activation of several signaling cascades. Disruption of this gene is associated with focal epilepsy and speech disorder with or without mental retardation. Alternative splicing results in multiple transcript variants.

Reference

Anti-NMDAR2A/GRIN2A Antibody被引用在1文献中。

Selected Validation Data



Western blot analysis of NMDAR2A/GRIN2A using anti-NMDAR2A/GRIN2A antibody (PB0323). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: Rat Brain tissue lysates,

Lane 2: Mouse Brain tissue lysates.

After electrophoresis, proteins were transferred to a membrane.

Then the membrane was incubated with rabbit anti-NMDAR2A/GRIN2A antigen affinity purified polyclonal antibody (PB0323) at a dilution of 1:1000 and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1197). A specific band was detected for NMDAR2A/GRIN2A at approximately 165 kDa. The expected band size for NMDAR2A/GRIN2A is at 165 kDa.