

Basic Information

Product Name	Anti-MMP9 Antibody
Gene Name	MMP9
Source	Rabbit
Clonality	Polyclonal
Isotype	IgG
Species Reactivity	mouse, rat
Tested Application	WB, IHC
Contents	500 ug/ml antibody with PBS, 0.02% NaN3, 1 mg/ml BSA and 50% glycerol.
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of mouse MMP-9, different from the related human sequence by thirteen amino acids, and from the related rat sequence by eight amino acids.
Concentration	500 ug/ml
Purification	Immunogen affinity purified.
Observed MW	78 kDa
Dilution Ratios	Western blot (WB): 1:500-2000 Immunohistochemistry (IHC): 1:50-400 (Boiling the paraffin sections in 10mM citrate buffer,pH6.0,or PH8.0 EDTA repair liquid for 20 mins is required for the staining of formalin/paraffin sections.) Optimal working dilutions must be determined by end user.

Storage

12 months from date of receipt, -20°C as supplied.

Background Information


Matrix metalloproteinase 9 (MMP-9), also known as 92 kDa type IV collagenase, 92 kDa gelatinase or gelatinase B (GELB), is an enzyme that in humans is encoded by the MMP9 gene. Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes. Most MMPs are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. The enzyme encoded by this gene degrades type IV and V collagens. Studies in rhesus monkeys suggest that the enzyme is involved in IL-8-induced mobilization of hematopoietic progenitor cells from bone marrow, and murine studies suggest a role in tumor-

associated tissue remodeling.

Reference

Anti-MMP9 Antibody被引用在60文献中。

Selected Validation Data



Western blot analysis of MMP9 using anti-MMP9 antibody (PB9669). The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: NRK whole cell lysates,

Lane 2: ANA-1 whole cell lysates,

Lane 3: HEPA whole cell lysates.

After electrophoresis, proteins were transferred to a membrane.

Then the membrane was incubated with rabbit anti-MMP9 antigen

affinity purified polyclonal antibody (PB9669) at a dilution of 1:1000

and probed with a goat anti-rabbit IgG-HRP secondary antibody

(Catalog # BA1054). The signal is developed using ECL Plus Western

Blotting Substrate (Catalog # AR1197). A specific band was detected

for MMP9 at approximately 78 kDa. The expected band size for

MMP9 is at 81 kDa.

Product datasheet

Anti-MMP9 Antibody

Catalog Number: **PB9669**


BOSTER[®]

antibody and ELISA experts

BOSTER BIOLOGICAL TECHNOLOGY

Building C21, 3rd to 5th Floors, Optics Valley Biopharmaceutical Accelerator,
East Lake High-Tech Development Zone, Wuhan.

Web: www.boster.com **Phone:** 027-67845390/1/2 **Email:** boster@boster.com



IHC analysis of MMP9 using anti-MMP9 antibody (PB9669).

MMP9 was detected in a paraffin-embedded section of mouse kidney tissue. Biotinylated goat anti-rabbit IgG was used as secondary antibody. The tissue section was incubated with rabbit anti-MMP9 Antibody (PB9669) at a dilution of 1:200 and developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB (Catalog # AR1027) as the chromogen.